AD-A214 713 NTATION PAGE

Form Approved OMB No. 0/04-0188

1. AGENCY USE ONLY (Leave blank)	2. REPORT DATE May 84	3. REPORT TYPE AN Final (5/	D DATES COVERED 15/82-5/14/83
4. TITLE AND SUBTITLE			5. FUNDING HUMBERS
FINAL SCIENTIFIC REPORT 15 MAY 1982-14 MAY 1983		-0171,	61102F
6. AUTHOR(S)	 		2304/A4
Frederick Bloom			230 1/11.1
7. PERFORMING ORGANIZATION NAME	(S) AND ADDRESS(ES)		8. PERFORMING ORGANIZATION
University of South Car Dept of Mathema tics & Columbia SC 29208	olina	APTIN	REPORT NUMBER
9. SPONSORING/MONITORING AGENC	NAME(S) AND ADDRESS	(ES)	10. SPONSORING/MONITORING
AFOSR bldg 410		•	AGENCY REPORT NUMBER
BAFB DC 20332-6448			AFOSR-81-0171
11. SUPPLEMENTARY NOTES		t v	
12a. DISTRIBUTION/AVAILABILITY STA	TEMENT	, , , , , , , , , , , , , , , , , , ,	12b. DISTRIBUTION CODE
DISTRIBU	TION STATEMENT A		
	d for public release; ibution Unlimited		
13. ABSTRACT (Maximum 200 words)			L
"Systems of nonlinear h electromagnetic theory,	yperbolic equation " "Shock formation " electromagnetic " systems associa	ons associated with on in inhomogeneou ic models," and "S ated with nonlinea	Shock wave formation for ar transmission line e report.
		1	DTIC
			NOV 29 1989
14. SUBJECT TERMS			15. NUMBER OF PAGES
			16. PRICE CODE
17 SECURITY CLASSIFICATION 18. OF REPORT	SECURITY CLASSIFICATION	N 19. SECURITY CLASSIF	CATION 20. LIMITATION OF ABSTRAC

unclassified

NSN 7540-01-280-5500

unclassified

Standard Form 298 (890104 Draft) frescribes by AMS 546, 239-18 299-81

Northern Illinois University DeKalb, Illinois 60115-2888

Department of Mathematical Sciences 815-75

May 14, 1984

APOSR TR. ×9-

TO: AFOSR

Bolling, AFD, DC 20332

FROM: Frederick Bloom, Professor

Department of Mathematical Sciences, NIU

RE: Final Scientific Report, AFOSR-81-0171, 15 May 1982 - 14 May 1983

During the period 15 May 1981 - 14 May 1982 the principal investigator was on leave at the University of Connecticut (Fall 1981) and the University of Maryland (Spring and Summer 1982); work completed during that period is summarized in the enclosed interium report covering that period. During the period 15 May 1982 - 14 May 1983 our efforts were centered on studying shock formation for plane waves propagating in nonlinear dielectric media and on the possible dissipative effects of anisotrophy, nonlinear conduction currents, and nonlinear dielectric relaxation. We also considered problems of singularity formation for electromagnetic wave propagation in nonlinear distributed parameter transmission lines. Appropriate summaries of our work in this area, some of which was completed after the principal investigator had accepted his present position at Northern Illinois University, may be found in the attached abstracts #4, 5, and 6.

:sec

Ful Bloom

Systems of Nonlinear Hyperbolic Equations
Associated with Problems of Classical
Electromagnetic Theory

Frederick Bloom

Department of Mathematical Sciences
Northern Illinois University
DeKalb, Illinois 60115

Abstract

The manifestation of a variety of nonlinear phenomena in electromagnetic theory is considered with specific emphasis on shock formation for plane waves propagating in nonlinear dielectric media and the possible dissipative effects of anisotrophy, nonlinear conduction currents, and nonlinear dielectric relaxation. Also considered are problems of singularity formation for electromagnetic wave propagation in nonlinear distributed parameter transmission lines.



Access	ion For	
NTIS	GRASI	
DTIC T		
Unsmoo		
វីម១៦រវ	teation.	
<u> </u>		
Ву		
Distr	(bution/	
Ava1	lability	Codes
	Avail a	ro\bn
Dist	Specia	11
1		
	}	
IN,	1	

Shock Formation in Inhomogeneous Quasilinear Systems Associated with Nonlinear Electromagnetic Models

Frederick Bloom

Department of Mathematical Sciences Northern Illinois University DeKalb, Illinois 60115

ABSTRACT

We consider the problem of the existence of smooth globally defined solutions to a quasilinear system of inhomogeneous hyperbolic equations; these equations are shown to model both the behavior of a distributed parameter non-linear transmission line, with voltage dependent capacitance and non-zero resistance and leakage conductance, and the interaction of a plane electromagnetic wave with a nonlinear dielectric half-space in the presence of a nonlinear conduction current. Specific conditions are exhibited, for both models, under which shocks form if the gradients of the initial data in each respective problem are, pointwise, sufficiently large. The analysis is based upon a study of the behavior of appropriate sets of Riemann invariants along their respective characteristic curves.

SHOCK NAVE FORMATION FOR INHOMOGENEOUS HYPERBOLIC SYSTEMS ASSOCIATED WITH NONLINEAR TRANSMISSION LINE PROBLEMS

Frederick Bloom
Department of Mathematical Sciences
Northern Illinois University
Dekalb, Illinois 60115

Abstraci

The problem of the existence of smooth globally defined solutions to a quasilinear system of inhomogenous hyperbolic equations modelling the behavior of a distributed parameter nonlinear transmission line is considered; the model assumes constant self-inductance, voltage dependent capacitance, and non-zero resistance and leakage conductance. Specific conditions are exhibited under which shocks form in solutions of the transmission line equations if at some point in the line the initial gradients of charge and current are sufficiently large.

Introduction

We consider a distributed parameter nonlinear transmission line with constant self-inductance L, current i, resistance R, and leakage conductance $\frac{1}{G}$ per unit length of the line; it is assumed that Q = Q(v) where v = v(x,t) is the voltage at a point in the line x units distance from an origin Q chosen in the line. The capacitance C is then given by $C \equiv C(v) = dQ(v)/dV$ and is, thus, also voltage dependent. The entire situation is depicted below:

$$\frac{\partial u_j}{\partial t} + \frac{\partial f_j}{\partial x} (\underline{u}) = g_{\underline{I}} (\underline{u}) \tag{3}$$

$$\underline{u} = \begin{pmatrix} 1 \\ Q \end{pmatrix}, \ \underline{f}(\underline{u}) = \begin{pmatrix} \frac{1}{L} V(Q) \\ \frac{1}{L} \end{pmatrix}, \ \underline{g}(\underline{u}) = -\begin{pmatrix} \frac{R}{L} & 1 \\ GV(Q) \end{pmatrix}$$
 (4)

and associated with (2) are the real characteristics in the x,t plane defined by $\frac{dx}{dt} = \frac{1}{\sqrt{L}} \sqrt{V^*(Q)}$.

Remarks.

(1) If G ≡ 0 then the system (2) reduces to

$$\begin{cases} Q_{,t} + 1, x = 0 \\ i_{,t} + 1/L(V(Q))_{,x} = -(R/L)1 \end{cases}$$
 (S.)

which is a special case of the damped quasilinear system

$$\begin{cases} \frac{\partial u}{\partial t} - \frac{\partial v}{\partial t} = 0 \\ \frac{\partial v}{\partial t} - \Gamma^*(u) \frac{\partial u}{\partial x} = -\alpha v \end{cases} \tag{6}$$

considered by Nishida [1] and Slemrod [2] for the care of associated periodic initial data $w(x,0) = w_0(x)$ $v(x,0) = v_0(x)$. It was shown in [1] that if |v(0)| > 0, |v(0)| > 0, and $||w_0||_{L^{\infty}}$, $||v_0||_{L^{\infty}}$.